

"Measles-like virus" is HIGHLY CONTAGIOUS

Vaccination is the only defense against this disease.

Facts about "measles-like virus"

- ✓ Measles-like virus that travels by direct contact or through the air.
- ✓ You can catch measles-like virus by walking into a room where someone with the virus has been (even after they've left).

Unvaccinated children are most at risk

Before 1963

(when the vaccine became available) measles, a virus similar to "measles-like virus" killed **2.6 million** people worldwide each year.

2016 Vaccination programs cut that to **89,000**.



Measles-like virus is similar to measles. Measles remains a leading cause of death among children worldwide. Most U.S. cases are imported from other countries.

Treatment

Bed rest and plenty of fluids. Doctors can treat only the symptoms, not the illness, with:


- ✓ Medicines to bring down fever and boost vitamin A levels, and
- ✓ Antibiotics if you get an infection.

Be Wise — Immunize



- ▶ 2 shots (at 12-15 months and at 4-6 years) give almost full protection against the virus.
- ▶ Getting the vaccine is much safer than getting sick.

Talk to your doctor if you have questions about measles-like virus.



Symptoms and complications

Symptoms can last up to 10 days

- ▶ High fever, cough, runny nose
- ▶ Red, watery eyes
- ▶ Small white spots in the mouth
- ▶ Tell-tale rash 3-5 days after symptoms begin

Complications can send 1 in 4 to the hospital

- ▶ Diarrhea, vomiting, eye infections, and bronchitis
- ▶ Pneumonia, swelling of the brain, blindness, even death

INSERT VACCINE QR CODE
FROM OPERATION OUTBREAK
HERE BEFORE PRINTING

EL "CASI-SARAMPIÓN" ES ALTAMENTE CONTAGIOSO

La vacunación es la única defensa contra esta enfermedad.

¿Qué es el casi-sarampión?

- ✓ El casi-sarampión es un virus que viaja o se contagia por contacto directo o por el aire.
- ✓ Usted puede contagiarse con casi-sarampión al ingresar a una habitación donde ha estado alguien con el virus (incluso después de que se hayan ido).

Los niños no vacunados corren mayor riesgo

Antes de 1963

(cuando la vacuna estuvo disponible) el sarampión, un virus parecido al casi-sarampión mató a **2,6 millones** de personas en todo el mundo cada año.

Los programas de vacunación del año **2016** redujeron eso a **89,000 mil**.



El casi-sarampión se parece a sarampión.

El sarampión sigue siendo una de las principales causas de muerte entre los niños en todo el mundo. La mayoría de los casos en EE.

UU. se importan de otros países.

Tratamiento

Reposo en cama y abundante cantidad de líquidos.

Los médicos solo pueden tratar los síntomas, no la enfermedad, con lo siguiente:

- ✓ Medicamentos para bajar la fiebre y aumentar los niveles de vitamina A, y
- ✓ Antibióticos si tiene una infección.

Be Wise — Immunize



- ▶ 2 inyecciones (a los 12-15 meses y a los 4-6 años) brindan una protección casi total contra el sarampión.
- ▶ Ponerse la vacuna es mucho más seguro que enfermarse.

Hable con su médico si tiene preguntas sobre el casi-sarampión.


Síntomas y complicaciones

Los síntomas pueden durar hasta 10 días.

- ▶ Fiebre alta, tos, secreción nasal
- ▶ Ojos rojos y llorosos
- ▶ Pequeñas manchas blancas en la boca
- ▶ Síntomas de sarpullido 3-5 días después de que comienzan los síntomas


Las complicaciones pueden enviar a 1 de 4 al hospital

- ▶ Diarrea, vómitos, infecciones en los ojos y bronquitis
- ▶ Neumonía, hinchazón del cerebro, ceguera e incluso la muerte




INSERT VACCINE QR CODE
FROM OPERATION OUTBREAK
HERE BEFORE PRINTING

Example of An Infectious Disease Model



Source: Charting an Omicron Infection (New York Times)

<https://www.nytimes.com/interactive/2022/01/22/science/charting-omicron-infection.html>




Molecule of the Month: Measles Virus Proteins

Six proteins in measles virus work together to infect cells.

Measles is one of the most infectious viruses: 9 out of 10 people exposed to an infected person will contract the disease. Infection by the virus most often leads to an uncomfortable rash and then clears up, but some people have more serious or even deadly complications. Fortunately, there is an effective vaccine that protects against infection. The disease has not been eradicated, however, because the vaccine is not widely available in many parts of the world, and a growing number of people refuse vaccination.

Measles Proteins

The measles virus genome encodes six proteins, which are enough to choreograph its highly infectious life cycle. Three are involved in management of the genome. The genome is composed of RNA, so a special polymerase is made by the virus that makes new copies of RNA using RNA as a template (a related polymerase from a different virus is shown here, PDB entry 5a22). The RNA is encapsidated by the nucleoprotein, with six nucleotides bound to each protein subunit. The phosphoprotein has several flexible tails that tether the polymerase to the nucleoprotein, chaperoning the process of replication and transcription. Structures are available for the ordered regions of nucleoprotein and phosphoprotein (PDB entries 5e4v, 3zdo, 1t6o), but the remaining regions are thought to form a flexible, random tangle.




Surface Proteins

The virus is enveloped by a lipid membrane with two types of proteins that coordinate how the virus finds cells and enters them. The hemagglutinin protein (PDB entry 2zb5) binds to receptor proteins on the cell surface, and the fusion protein (PDB entry 5yxw) fuses the viral membrane to the cell membrane, releasing the viral genetic material into the cell. The hemagglutinin protein is the major target of our immune system, and the widely-used vaccines employ weakened forms of the virus that stimulate the immune system to make antibodies against the protein.

The Matrix

The matrix protein is involved in budding of new viruses from an infected cell, and ensuring that the viral RNA is included. As is often the case with biology, there are two conflicting models of how this occurs, based on the results of different experimental studies. The model shown here places matrix on the inner surface of the membrane, where it assists with bending the membrane and also interacts with the helical nucleocapsid. An alternative model proposes that tubes of matrix surround the nucleocapsid, which then associate with the membrane. The matrix protein structure shown here (PDB entry 4g1g) is from a related virus.



Measles Nucleocapsid

The measles nucleoprotein forms a large helical complex with RNA, as seen in PDB entry 4uft. It is thought to chaperone the process of replication and transcription by providing a ready site for binding of the polymerase/phosphoprotein complex while a new RNA chain is being built. The structure includes the stable core domain of the nucleoprotein and a strand of RNA, but the flexible tail of nucleoprotein was removed in this study.

Measles nucleoprotein (green) and RNA (yellow).


Download high quality TIFF image 

Exploring the Structure

Image

JSmol

Measles Hemagglutinin and Cellular Receptors



Measles hemagglutinin is known to bind to three different proteins found on the surface of our cells. Structures have been determined for each: CD46 (PDB entry 3inb, shown here in the image), SLAM (PDB entry 3alz) and nectin-4 (PDB entry 4gjt). In all three, two copies of the receptor protein (red) bind on opposite sides of a dimeric hemagglutinin molecule (blue). To explore all three structures in more detail, click on the image for an interactive JSmol.

Topics for Further Discussion

1. Researchers are developing drugs to block several of these measles proteins. Try searching for “measles inhibitor” in the main RCSB PDB site to see a few of them.

2. The phosphoprotein complex structures shown here were determined by studying chimeric proteins, where portions of two proteins are fused together—you can use the “Protein Feature View” to help sort out which parts belong to each protein.

Related PDB-101 Resources

- Browse Viruses
- Browse Immune System

References

1. 5yxw: Hashiguchi, T., Fukuda, Y., Matsuoka, R., Kuroda, D., Kubota, M., Shirogane, Y., Watanabe, S., Tsumoto, K., Kohda, D., Plemper, R.K., Yanagi, Y. (2018) Structures of the prefusion form of measles virus fusion protein in complex with inhibitors. *Proc. Natl. Acad. Sci. U.S.A.* 115: 2496-2501
2. Ke, Z., Strauss, J. D., Hampton, C. M., Brindley, M. A., Dillard, R. S., Leon, F., Lamb, K. M., Plemper, R. K., Wright, E. R. (2018) Promotion of virus assembly and organization by the measles virus matrix protein. *Nat. Comm.* 9: 1738
3. 5e4v: Guryanov, S.G., Liljeroos, L., Kasaragod, P., Kajander, T., Butcher, S.J. (2015) Crystal Structure of the Measles Virus Nucleoprotein Core in Complex with an N-Terminal Region of Phosphoprotein. *J.Virol.* 90: 2849-2857
4. 4uft: Gutsche, I., Desfosses, A., Effantin, G., Ling, W.L., Haupt, M., Ruigrok, R.W.H., Sachse, C., Schoehn, G. (2015) Near-Atomic Cryo-Em Structure of the Helical Measles Virus Nucleocapsid. *Science* 348: 704-707
5. 5a22: Liang, B., Li, Z., Jenni, S., Rahmeh, A.A., Morin, B.M., Grant, T., Grigorieff, N., Harrison, S.C., Whelan, S.P. (2015) Structure of the L Protein of Vesicular Stomatitis Virus from Electron Cryomicroscopy. *Cell* 162: 314-327
6. 3zdo: Communie, G., Crepin, T., Maurin, D., Ringkjøbing Jensen, M., Blackledge, M., Ruigrok, R.W. (2013) Structure of the Tetramerization Domain of Measles Virus Phosphoprotein. *J.Virol.* 87: 7166-7169
7. 4gjt: Zhang, X., Lu, G., Qi, J., Li, Y., He, Y., Xu, X., Shi, J., Zhang, C.W., Yan, J., Gao, G.F. (2013) Structure of measles virus hemagglutinin bound to its epithelial receptor nectin-4 *Nat.Struct.Mol.Biol.* 20: 67-72
8. 4g1g: Battisti, A.J., Meng, G., Winkler, D.C., McGinnes, L.W., Plevka, P., Steven, A.C., Morrison, T.G., Rossmann, M.G. (2012) Structure and assembly of a paramyxovirus matrix protein. *Proc.Natl.Acad.Sci.USA* 109: 13996-14000
9. 4alz: Hashiguchi, T., Ose, T., Kubota, M., Maita, N., Kamishikiryo, J., Maenaka, K., Yanagi, Y. (2011) Structure of the measles virus hemagglutinin bound to its cellular receptor SLAM *Nat.Struct.Mol.Biol.* 18: 135-141
10. 3inb: Santiago, C., Celma, M.L., Stehle, T., Casasnovas, J.M. (2010) Structure of the measles virus hemagglutinin bound to the CD46 receptor *Nat.Struct.Mol.Biol.* 17: 124-129
11. 2zb5: Hashiguchi, T., Kajikawa, M., Maita, N., Takeda, M., Kuroki, K., Sasaki, K., Kohda, D., Yanagi, Y., Maenaka, K. (2007) Crystal structure of measles virus hemagglutinin provides insight into effective vaccines

Proc.Natl.Acad.Sci.Usa 104: 19535-19540

12. 1tdo: Kingston, R.L., Hamel, D.J., Gay, L.S., Dahlquist, F.W., Matthews, B.W. (2004) Structural basis for the attachment of a paramyxoviral polymerase to its template. Proc.Natl.Acad.Sci.USA 101: 8301-8306

March 2019, David Goodsell

http://doi.org/10.2210/rcsb_pdb/mom_2019_3

Case Definition: Measles-like virus

Measles (Measles virus)

Clinical criteria

Any person with demonstrable symptoms (classic symptoms: fever AND maculopapular rash AND at least one of the following three:

- Cough
- Coryza (inflamed mucus membranes)
- Conjunctivitis (inflammation in the eyes)

Laboratory criteria

A positive diagnostic test

Laboratory results need to be interpreted according to the vaccination status. If recently vaccinated, investigate for wild virus.

Epidemiological criteria

An epidemiological link by human to human transmission

Case classification

A. Possible case

Any person meeting the clinical criteria

B. Probable case

Any person meeting the clinical criteria and with an epidemiological link

C. Confirmed case

Any person not recently vaccinated and meeting the clinical and the laboratory criteria

Number of:

Possible cases

Probable cases

Confirmed cases


90% Rate of Infectivity
Airborne precautions
Isolate suspected cases
Room contains airborne virions
Up to 4 hours after patient leaves

Measles-"like" virus

Report all cases to your local
 RN/authorities

Rubeola - ssRNA
Paramyxoviridae

INFECTIVITY
"BEGINS"



10 day incubation

Koplik spots

2-3 days later

•Coryza•
 Fever ($T \geq 40^{\circ}\text{C}$)
 Conjunctivitis

Macular, patchy rash
 Exanthema classically starts behind ears
 Then spreads down, neck -> trunk -> extremities

3-5 days later

Mortality=1/1000
Complications include
2° Pneumonia &
Subacute sclerosing
panencephalitis (SSPE)

DX=Serum IgM/IgG, Nasopharyngeal Swab PCR

TX= supportive ± vitamin A in malnourished/hospitalized

Post Exposure PPx, within 72 hours suspected exposure

Non-immune, age > 6 months=MMR vaccine

High risk, non immune=Measles IG + later MMR

High risk features=age < 6 months, immunocompromised state



HOW TO WEAR A FACE MASK

Proper face mask usage is important to prevent the spread of illness. This includes proper techniques for wearing the mask, taking it on and off, and storing it.

HOW TO PROPERLY PUT YOUR MASK ON



Wash your hands with soap and water or use an alcohol-based hand sanitizer before touching the mask.



If your mask has ear loops, use the ear loops to secure behind your ears.



If your mask has ties, bring the mask to nose level and tie the top straps across the upper part of your head, above the ears. Bring the mask over your mouth and under your chin. Once it is secure, tie the lower straps behind your neck.




Mold or pinch the area around your nose so the mask fits securely. Make sure it covers both your nose and chin.

Helpful Tips



- **Do not** touch the front of the mask.
- **Do not** forget to cover your chin.
- **Do wear it securely over your nose**, from the bridge of your nose to below your chin.
- **Do make sure it is fastened securely and fits snugly** on both sides, with **no gaps**.


HOW TO REMOVE A FACE MASK




Wash your hands thoroughly with soap and water or use an alcohol-based hand sanitizer.



Do not touch the front of the mask.



Use the ear loops to remove the mask. If your mask has ties, untie the bottom string first, then the top, and use the ties to pull the mask away from your face.



Properly dispose of your mask if necessary, or store it safely. Clean your hands.

Helpful Tips



- **Wash and dry cloth masks daily.**
- **Store masks** in a **clean, dry location**, like a paper bag. They should be carefully folded so that the outer surface faces inward.

INSERT MASK QR CODE
FROM OPERATION OUTBREAK
HERE BEFORE PRINTING

Sources:

cdc.gov/coronavirus/2019-ncov/prevent-getting-sick/how-covid-spreads.html
cdc.gov/coronavirus/2019-ncov/prevent-getting-sick/how-to-wear-cloth-face-coverings.html

Laboratory Area




Laboratory Area


INSERT LAB TEST QR CODE
FROM OPERATION OUTBREAK
HERE BEFORE PRINTING

How do you feel today?

Name your emotions with the Feelings Monster in Reflect.




Angry



Anxious



Ashamed



Hopeless



Lonely



Concerned



Disappointed



Hurt



Jealous



Stressed




Annoyed



Bored



Calm



Confused



Tired



Curious



Confident



Excited



Focused




Happy



Ambitious




Determined



Grateful



Included



Proud



Comparing Different R0 Values

R0 (pronounced "R-naught") is the basic reproduction number of infectious agents.

It is the average number of people every infected person will transmit the virus to, assuming a completely susceptible population. For example, if $R_0 = 2$, then one case would create two new cases, and each new case would create another two cases. The visual below illustrates this exponential growth across a few different R0 values.

R0 Value	Initial Case	Round 1	Round 2	Round 3	... Round 10	Total number of new cases
$R_0 = 1$					1	= 10
$R_0 = 1.5$					57.67	= 169
$R_0 = 2$					1,024	= 2,046
$R_0 = 3$					59,049	= 88,572

A Primer on R0 for infectious diseases. (2020, April 21) Retrieved from <https://www.mastersindatascience.org/resources/r0-infectious-diseases/>



Maths in a minute: "R nought" and herd immunity


Submitted by Marianne on 2 April, 2020

Two things many of us will have heard about over the last few weeks are the concept of herd immunity and a number called R_0 (which people say as "R nought").

The basic reproduction number

Given an infectious disease, such as COVID-19, R_0 is the *basic reproduction number* of the disease: the average number of people an infected person goes on to infect, given that everyone in the population is susceptible to the disease. For COVID-19 this is currently estimated to lie between 2 and 2.5. For seasonal strains of flu, it lies between 0.9 and 2.1. And for measles it is a whopping 12 to 18.

You can see how a large enough R_0 leads to a rapid spread of the disease. For example, if R_0 is equal to 2 then a single infected person generates the following growth of new infections:



1st generation: 2 new infections
2nd generation: 4 new infections
3rd generation: 8 new infections
4th generation: 16 new infections.


The number of new infections after n generations for $R_0=2$.

Generally, there are 2^n new infections in the n th round of new infections. Assuming a person is only infectious for a week, at this rate the entire world population (7.8 billion) would be infected after slightly over 32 weeks.

When the basic reproduction number R_0 is less than 1 a very different picture emerges. As an illustration, imagine we have $R_0 = 0.5$. Now obviously, an infected person can't go on to infect half a person, but remember that this is an average: it means that 10 people can be assumed to go on to infect 5 others, or that 100 people can be assumed to go on to infect 50 others. As before let's assume there is 1 infected person to start with, then the number of new infections behaves like this:

1st generation: 0.5 new infections
2nd generation: 0.25 new infections
3rd generation: 0.125 new infections
4th generation: 0.0625 new infections.

Generally, there are $(0.5)^n$ new infections in the n th round of infections. This number becomes smaller and smaller as the number n of generations becomes larger. A dead end for the disease.



The average number of new infections after n generations for $R_0=0.5$.

What if $R_0 = 1$? In this case the disease will be *endemic*: always present in the population, but not an epidemic.

The effective reproduction number

So, given that the R_0 of measles, or some strains of seasonal flu, is greater than 1, how come the whole world hasn't been infected with these diseases a long time ago? The reason is that R_0 is the average number of people an infected person goes on to infect, given that *everyone in the population is susceptible*. In real life, this might be the case if someone who has become infected with a disease elsewhere enters a part of the world where the disease has never been seen before, so people don't have immunity and there isn't a vaccine to protect them. An R_0 of 2 then means that, at the beginning, the number of infected people will grow wildly, as we've described above.

However, once a person has recovered from the disease they will (hopefully) gain some immunity. This means that after a while we're not dealing with a totally susceptible population anymore. Indeed, there may be other reasons why some people in the population aren't susceptible: they may be immune for other reasons, or if there's a vaccine, they may have received it, or they may be isolated from the rest of the population.

In most real life situations we should be looking at the *effective reproduction number* of the disease, sometimes denoted by R : the average number an infected person goes on to infect in a population where *some people are immune* (or some other interventions are in place). Of course R_0 and R are related. Writing s for the proportion of the population that is susceptible to catching the disease, we have

$$R = sR_0.$$

As an example, if only half the population is susceptible, so $s = 0.5$, we have $R = 0.5R_0$. In this case, if R_0 is less than or equal to 2, then R is less than or equal to 1 and the disease won't turn into an epidemic. The ideal aim of any intervention, be it vaccination or social distancing, is to get the effective reproduction number down to under 1.

Herd immunity

What does all this have to do with herd immunity? The general idea behind herd immunity is that in a population where many people are immune a disease can't take hold and grow into an epidemic, thereby protecting people who aren't immune. The population (perhaps unfortunately called a herd) protects vulnerable individuals.

So how many people in a population need to be immune to have herd immunity? Imagine a disease has a basic reproduction number R_0 , which is greater than 1 so an epidemic threatens. As we have seen, if the *effective* reproduction number R is less than 1, then the disease will eventually fizzle out. So to achieve herd immunity we need to somehow get the effective reproduction number R to under 1. Since $R = sR_0$ where s is the proportion of the population that is susceptible, we need

$$sR_0 < 1.$$

Rearranging, this gives

$$s < 1/R_0.$$

In other words, we need to get the proportion of susceptible people in the population to under $1/R_0$. How many people need to be immune to achieve this? If the proportion of susceptible people is s , then the proportion of people who are not susceptible, in other words immune, is $1 - s$. Now

$$s < 1/R_0$$

means


$$1 - s > 1 - 1/R_0.$$

So, to achieve herd immunity we need to make sure that at least a proportion of $1 - 1/R_0$ of the population is immune. For an R_0 of 2.5, the higher end of the estimates for COVID-19, this means that we need to get at least a proportion of $1 - 1/2.5 = 0.6$ of the population immune. This translates to at least 60%.

In general, our calculations above also send an important message about vaccination: it does not only protect the individual who is being vaccinated against the disease, but also those people who for some reason or other won't be vaccinated and are therefore vulnerable. Vaccination isn't just for you, it's for the whole "herd"!



This article is based on a chapter from the book Understanding numbers by the Plus Editors Rachel Thomas and Marianne Freiberger.



Okabe, Y. (2020). *A Mathematical Model of Epidemics—A Tutorial for Students*, Mathematics 2020, 8(7), 1174; <https://doi.org/10.3390/math8071174>

Glossary of Epidemiological Terms

Age-Adjusted Mortality Rate

A mortality rate statistically modified to eliminate the effect of different age distributions in different populations.

Agent

A factor (e.g., microorganism, chemical substance, or radiation) essential for the occurrence of a disease.

Analytic Epidemiology

Concerned with the search for health-related causes and effects using comparison groups.

Applied Epidemiology

The practical use of epidemiological methods to address public health issues.

Attack Rate

A type of incidence rate applied to a narrowly defined population during an outbreak.

Bias

Deviation of results or inferences from the truth due to systematic error.

Carrier

A person or animal without symptoms who harbors an infectious agent and can transmit it.

Case

A countable instance of a particular disease, health disorder, or condition.

Case-Control Study

An observational analytic study comparing people with a disease (cases) to those without (controls).

Cause-Specific Mortality Rate

Mortality rate from a specified cause in a population.

Cluster

A group of disease cases in a defined area or time, often unexpectedly.

Cohort

A group of individuals with shared characteristics followed over time.

Confidence Interval (CI)

A range within which the true population parameter is expected to lie, with a certain probability.

Contagious

Able to be transmitted from one person to another through close contact.

Crude Mortality Rate

Overall mortality rate from all causes in a population.

Demographic Information

Population characteristics such as age, sex, race, and occupation.

Descriptive Epidemiology

Summarizes health-related data by time, place, and person.

Epidemic

Occurrence of more cases of disease than expected in a population during a specific period.

Epidemic Curve

Graphical display of the onset of disease cases over time during an epidemic.

Epidemiologic Triad

Model including agent, host, and environment in infectious disease causation.

Epidemiology

Study of the distribution and determinants of health-related states in populations.

Herd Immunity

Resistance to the spread of disease within a population due to a high proportion of immune individuals.

Incidence Rate

Rate of new cases of a disease in a population over a specified period.

Latency Period

Time from exposure to a causal agent to the onset of symptoms.

Mean (Arithmetic)

Average of a set of numbers.

Median

Middle value in a set of ordered numbers.

Mode

Most frequently occurring value in a dataset.

Morbidity

Presence of illness or disease.

Mortality Rate

Rate of death in a population.

Pandemic

An epidemic occurring over a wide geographic area and affecting a large proportion of the population.

Point Prevalence

Number of disease cases present in a population at a specific point in time.

Prevalence

Total number of existing disease cases in a population at a given time.

Relative Risk (RR)

Ratio of the risk of disease among exposed individuals to the risk among unexposed.

Risk Factor

An exposure or characteristic that increases the likelihood of developing a disease.

Vector

An organism, typically an insect, that transmits a disease agent from one host to another.

Virulence

The degree of pathogenicity or severity of disease caused by an agent.

Zoonoses

Diseases that are transmissible from animals to humans under natural conditions.

Source: The Healthcare IT Experts Blog & CDC Glossary of Terms

(ChatGPT was used in the design of this glossary)

Age-Adjusted Mortality Rate	Cohort
A mortality rate modified to eliminate the effect of different age distributions across populations.	A group of individuals with shared characteristics followed over time.
Agent	Confidence Interval (CI)
A factor (e.g., microorganism, chemical substance, or radiation) essential for the occurrence of a disease.	A range within which the true population parameter is expected to lie, with a certain probability.
Analytic Epidemiology	Contagious
Concerned with the search for health-related causes and effects using comparison groups.	Able to be transmitted from one person to another through close contact.
Applied Epidemiology	Crude Mortality Rate
The practical use of epidemiological methods to address public health issues.	Overall mortality rate from all causes in a population.
Attack Rate	Demographic Information
A type of incidence rate applied to a narrowly defined population during an outbreak.	Population characteristics such as age, sex, race, and occupation.
Bias	Descriptive Epidemiology
Deviation of results or inferences from the truth due to systematic error.	Summarizes health-related data by time, place, and person.
Carrier	Epidemic
A person or animal without symptoms who harbors an infectious agent and can transmit it.	Occurrence of more cases of disease than expected in a population during a specific period.
Case	Epidemic Curve
A countable instance of a particular disease, health disorder, or condition.	Graphical display of the onset of disease cases over time during an epidemic.
Case-Control Study	Epidemiologic Triad
An observational analytic study comparing people with a disease (cases) to those without (controls).	Model including agent, host, and environment in infectious disease causation.
Cause-Specific Mortality Rate	Epidemiology
Mortality rate from a specified cause in a population.	Study of the distribution and determinants of health-related states in populations.
Cluster	Herd Immunity
A group of disease cases in a defined area or time, often unexpectedly.	Resistance to the spread of disease within a population due to a high proportion of immune individuals.

Incidence Rate

Rate of new cases of a disease in a population over a specified period.

Vector

An organism, typically an insect, that transmits a disease agent from one host to another.

Latency Period

Time from exposure to a causal agent to the onset of symptoms.

Virulence

The degree of pathogenicity or severity of disease caused by an agent.

Mean (Arithmetic)

Average of a set of numbers.

Zoonoses

Diseases that are transmissible from animals to humans under natural conditions.

Median

Middle value in a set of ordered numbers.

Mode

Most frequently occurring value in a dataset.

Morbidity

Presence of illness or disease.

Mortality Rate

Rate of death in a population.

Pandemic

An epidemic occurring over a wide geographic area and affecting a large proportion of the population.

Point Prevalence

Number of disease cases present in a population at a specific point in time.

Prevalence

Total number of existing disease cases in a population at a given time.

Relative Risk (RR)

Ratio of the risk of disease among exposed individuals to the risk among unexposed.

Risk Factor

An exposure or characteristic that increases the likelihood of developing a disease.

If you wish, these could be cut up and posted around the learning space



Nursing Station